

BASIC DE-WORMING GUIDELINES FOR MARES & FOALS

- De-worm pregnant mares as usual for the duration of the pregnancy. Include rotational de-wormers.
- De-worm mares 2-4 weeks prior to foaling.
- De-worm foal at 4 weeks of age. De-worm the mare at the same time.
- De-worm the foal every 4 weeks until 6 months old, then every 6 weeks until one year of age, then every 8 weeks from one year of age.
- De-worm mares every 6-8 weeks.

Reduce worm infestations by:

- Removing manure daily from stables, paddocks and yards.
- When hand feeding horses keep feed off the ground in feed bins or tyre feeders.
- Keep stocking rates as low as possible. Cross graze pasture with cattle & sheep.
- If possible, group horses in age categories. Put young stock in the least contaminated paddocks.
- Drench all horses on the property at the same time.
- Quarantine new horses & drench before mixing with other horses.

This is a guide only. De-worming programs vary depending on management strategies, the environment & number of stock. Please ask one of our vets for advice for your worm control program.



Consultations are by appointment.

Orange Veterinary Hospital

57 Molong Rd,
ORANGE, NSW 2800
PH: 02 63618388

Molong Veterinary Clinic

89 Bank St,
MOLONG, NSW 2866
PH: 02 63669021

OPENING HOURS:

MONDAY-FRIDAY 8:30AM TO 5:30PM
SATURDAY 8:30AM-12PM
24 HOUR EMERGENCY SERVICE



Vaccinating the Pregnant Mare and Foal



57 Molong Road
ORANGE NSW 2800
Phone: 02 6361 8388

Email: orangevet@orangevet.com.au
Website: www.orangevet.com.au
www.facebook.com/orangevethospital

PREGNANT MARE VACCINATIONS

Fully Vaccinated Mares:

Tetanus/Strangles - Previously vaccinated (up-to-date) mares require a single Equivac 2 in 1 booster 4-6 weeks prior to foaling.

Salmonella - Previously vaccinated mares (up-to-date) require a single Salmonella booster 4-6 weeks prior to foaling.

Equine Herpes Virus (EHV) - All pregnant mares should receive a Duvaxyn EHV booster at the beginning of the 5th, 7th & 9th month of pregnancy to help prevent abortion caused by EHV.

Uncertain Vaccination History:

Tetanus/Strangles - Unvaccinated mares require a Equivac 2 in 1 booster, repeated after 4 weeks. Ideally a booster vaccination should also be given 4-6 weeks prior to foaling.

Salmonella - Unvaccinated mares require a Salmonella vaccination, repeated at 4-6 weeks. Ideally a booster vaccination should also be given 4-6 weeks prior to foaling.

Equine Herpes Virus (EHV) - Vaccinate pregnant mares at the beginning of the 5th, 7th & 9th month of pregnancy.

FOAL VACCINATIONS

It is advisable to start foal vaccinations at 12 weeks of age.

First Vaccination:

(12 weeks old)

Equivac 2 in 1

(Tetanus/Strangles)

Salmonella

Duvaxyn EHV - If high risk of respiratory disease.

2 Weeks Later:

(14 weeks old)

Equivac S - (Strangles)

2 Weeks Later:

(16 weeks old)

Equivac 2 in 1

(Tetanus/Strangles)

Salmonella

4 Weeks Later:

(20 weeks old)

Duvaxyn (EHV)

4 Weeks Later:

(24 weeks)

Duvaxyn (EHV) - Booster every 6 months

Hendra Virus Vaccination can be given from 4 months of age– Two doses, three weeks apart followed by periodic boosters.

***Booster vaccinations for Strangles and EHV are recommended every 6 months for maximum protection.**

***Tetanus vaccination provides excellent immunity to the disease and is recommended annually.**

***Strangles, Salmonella & Equine Herpes Virus vaccinations help prevent these diseases & reduce the severity & duration of the disease, however total protection is not provided.**

WHY VACCINATE?

***These diseases can be severe & fatal**

***These diseases are reasonably common**

***Protection following vaccination is generally excellent provided regular boosters are given.**

***Vaccination is very inexpensive when compared to the cost of attempting to treat these diseases & the value of the horse.**

Note: Vaccination programs will differ for individual needs & situations. These are the recommendations for a complete vaccination program to protect against Tetanus, Strangles, Equine Herpes Virus and Salmonella in a stud situation. Please consult one of our vets for advice on a vaccination program and management strategies to suit your individual needs and environment.

